# **Original Article**

Nasrin Banu Rupani (MD) <sup>1</sup> Morteza Alijanpour (MD) <sup>2\*</sup> Kazem Babazadeh (MD) <sup>2</sup> Karimollah Hajian-Tilaki (PhD) <sup>3</sup> Fatemeh Moadabdoost (BSc) <sup>4</sup>

- 1. Student Research Committee, School of Medicine, Babol University of Medical Sciences, Babol, Iran 2. Non-Communicable Pediatric Diseases Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran 3. Babol University of Medical Sciences, Babol, Iran 4. The Clinical Research Development Unit of Amirkola Children's Hospital, Babol University of Medical Sciences, Babol, Iran
- \* Correspondence:
  Morteza Alijanpour, NonCommunicable Pediatric Diseases
  Research Center, Health Research
  Institute, Babol University of
  Medical Sciences, Babol, Iran

E-mail: m.alijanpour@yahoo.com Tel: 0098 1132354951 Fax: 0098 1132354951

Received: 11 Jun 2018 Revised: 16 Aug 2018 Accepted: 28 Aug 2018

# Effect of levothyroxine on cardiac function in children with subclinical hypothyroidism: A quasi-experimental study

## **Abstract**

**Background:** Because of the importance of thyroid function and its effect on different organs, such as the heart, this study was aimed to evaluate the effect of levothyroxine on cardiac function in children with subclinical hypothyroidism.

*Methods:* This quasi-experimental study was performed on children aged 4-12 years old with diagnosis of subclinical hypothyroidism in Amirkola Children's Hospital during 2018-2019. Cardiac functional parameters, including ejection fraction (EF), fractional shortening (FS), ratio of early filling velocity to early diastolic mitral annulus velocity (E/E'), myocardial performance index (MPI), left ventricular end-diastolic diameter (LVEDD), and left ventricular end-systolic diameter (LVESD), were measured by echocardiography at baseline and 6 months after levothyroxine treatment.

**Results:** Out of the 30 subjects, 19 (63.3%) were boys and 11 (36.7%) were girls. The mean age was  $6.60\pm2.13$  years old. The mean EF index was  $63.13\pm3.01$  percent before treatment, which increased to  $69.07\pm4.50$  percent after treatment (p<0.001). Also, the mean FS was  $31.83\pm1.62$  percent before treatment, which improved to  $35.10\pm1.13$  percent after treatment (p<0.001). The mean MPI was  $0.28\pm0.02$  before treatment, which increased to  $0.33\pm0.03$  after treatment (p<0.001). On the other hand, no significant difference was found in the mean E/E' before and after treatment. The mean LVEDD decreased from  $3.47\pm0.46$  cm before treatment to  $3.05\pm0.40$  cm after treatment (p<0.001), whereas LVESD non-significantly decreased after treatment versus before treatment.

*Conclusion:* The results showed that treatment with levothyroxine may improve cardiac functional parameters in children with subclinical hypothyroidism.

Keywords: Hypothyroidism, Child, Cardiac function, Levothyroxine

#### Citation:

Rupani NB, Alijanpour M, Babazadeh K, Hajian-Tilaki K, Moadabdoost F. Effect of levothyroxine on cardiac function in children with subclinical hypothyroidism: A quasi-experimental study. Caspian J Intern Med 2019; 10(3): 332-338.

Subclinical hypothyroidism is a branch of hypothyroidism which is characterized by increased thyroid stimulating hormone (TSH) and normal thyroxine (T4) (1). The prevalence of subclinical hyperthyroidism in pediatrics is about 2% and its incidence is more in girls than in boys (2, 3). Some of the complications of subclinical hypothyroidism include changing lipid profiles, increase blood coagulation and changing cardiac function especially left ventricle (4, 5). Cardiovascular system is sensitive to thyroid hormone activity. Although a wide spectrum of cardiac disorders has been identified in patients with thyroid dysfunctions, cardiac involvement in patients with subclinical hypothyroidism has been discussed over the recent two-three decades. Effects of subclinical hypothyroidism on cardiovascular system are directly related to severity of thyroid hormone deficiency. Subclinical hypothyroidism causes increase in systemic vascular resistance, diastolic dysfunction and decrease in preload of heart (6, 7). It can also lead to heart failure and coronary artery lesions (8).

There is conflicting evidence on the effect of subclinical hypothyroidism on decrease in systolic functions. Some papers stated that subclinical hypothyroidism can decrease systolic function (9, 10), but some others are not compatible with this result (11, 12). Thyroid hormones can affect cardiovascular system through transcriptional regulation and regulatory proteins (13, 14). In addition, hypothyroidism can contribute in chronic inflammations and tissue changes (such as dehydration and collagen alternation), and also hemodynamic changes by influence on arterial smooth muscle (15, 16). Cardiovascular complications of subclinical hypothyroidism can be seen in children and adults.

Some articles stated that treatment with levothyroxine was associated with positive effects on cardiac function in adults with subclinical hypothyroidism suffering from cardiovascular complications (17-20). However, number of studies assessing the effects of levothyroxine on cardiac complications in pediatric population is limited. Considering the importance of proper thyroid function and its effect on different organs, such as heart which has a vital role in correct function of various organs and in improving the growth and development of children, it would be clinically helpful to find a therapeutic option in pediatric patients with subclinical hypothyroidism to treat and prevent them from potential cardiac complications in the future. The purpose of the present study was to evaluate the effects of treatment with levothyroxine on cardiac function in the pediatric patients with subclinical hypothyroidism.

## **Methods**

This open-label quasi-experimental study was performed in 2018-2019 on children aged 4-12 years old, who were referred to the endocrinology clinic of the Amirkola Children's Hospital with the diagnosis of subclinical hypothyroidism, or the children who came for growth evaluation and were diagnosed as subclinical hypothyroidism by laboratory tests (elevated TSH and normal T4). Diagnosis of subclinical hypothyroidism was done by pediatric endocrinologist.

The exclusion criteria were as follows:

- 1. Being overweight (body mass index [BMI, kg/m²] between 85<sup>th</sup> and 95<sup>th</sup> percentile), obesity (BMI higher than 95<sup>th</sup> percentile), or underweight (BMI lower than 5<sup>th</sup> percentile) (21).
- 2. Hypertension (higher than 95<sup>th</sup> percentile for age, sex and height) (21)

- 3. Previous use of levothyroxine, anticonvulsants
- 4. Precocious puberty (22, 23)
- 5. Pathologic bradycardia or tachycardia (24)
- 6. Congenital or acquired heart disease and under medical treatment
- 7. Liver or kidney disease
- 8. Hyperlipidemia
- 9. Diabetes
- 10. Infectious diseases

All subjects were initially examined for thyroid, maturity stage, hypertension and pulse rate and their height and weight were recorded. The examination was conducted by the pediatric resident under supervision of the pediatric endocrinologist. Height and weight of the patients were evaluated and recorded by BALAS equipment (Nikan Andishan Nozhan Co, Tehran, Iran) with measurement error of 0.2 grams and then BMI was calculated using formula of weight (kilograms) divided by height (square meters). Blood pressure was assessed in the prone and rest position using the sphygmomanometer Heine (made in Germany). To conduct thyroid tests, 2 cc blood was collected after 8-10 hours fasting and then levels of TSH and T4 were measured using enzyme-linked immunosorbent assay (ELISA) kits (Pishtaz Teb, Tehran, Iran) in the laboratory of Amirkola Children's Hospital. Normal range for THS was 0.5-5.5 mIu/l and for T4 was 5.5-12.8 µg/dl (prepubertal children: 3-10 years old) or 4.2-13µg/dl (pubertal children >10 years old) (24).

After approving subclinical hypothyroidism, the subjects underwent echocardiography for the evaluation of cardiac function, using a Medison Accuvix V10 ultrasound equipment (Samsung Medison, Seoul, South Korea) by the pediatric cardiologist. Study outcomes and parameters of cardiac function assessed were as follows:

- Left ventricular end-systolic diameter (LVESD): Normal value depends on age, including  $1.9\pm0.3$  (1-5 years old),  $2.4\pm0.3$  (6-9 years old),  $2.7\pm0.3$  (10-13 years old) (25).
- Left ventricular end-diastolic diameter (LVEDD): Normal value depends on age, including 3.1±0.4 (1-5 years old), 3.9±0.4 (6-9 years old), 4.3±0.4 (10-13 years old) (25).
- Fractional shortening: It is defined as percentage of changes of left ventricular dimensions between end-diastole and end-systole (25). It is measured using the following formula and the normal range is between 28% and 38%: (LVEDD-LVESD)/LVEDD×100.
- Ejection fraction (EF): It is a volume parameter showing changes of left ventricular volume between end-diastole and

end-systole (25). It is measured using the following formula and the normal range is between 54% and 75%: (left ventricular end-diastolic volume [LVEDD]- left ventricular end-systolic volume [LVESD])/LVEDD×100.

- Ratio of early filling velocity to early diastolic mitral annulus velocity (E/E'): It is a strong predictor of post-acute myocardial infarction complications, hypertensive cardiac diseases, severe secondary mitral insufficiency, end stage renal disease, atrial fibrillation and cardiomyopathies (26). Normal value depends on age, including 8.1±1.8 (1-5 years old), 7.7±1.6 (6-9 years old), 6.6±1.4 (10-13 years old).
- •Myocardial performance index (MPI) or Tei index: It includes both systolic and diastolic time intervals to assess the global cardiac dysfunction (26). It is measured using the following formula and the normal value is  $0.35\pm0.03$  for left ventricle: (Isovolumic relaxation time+Isovolumic contraction time)/ Ejection time.

Patients with abnormal range of the mentioned parameters and clinical symptoms of heart failure, such as fatigue, anorexia, cough, dyspnea, sweating, abdominal pain, edema, hepatomegaly, orthopnea, rale in auscultation, and so on, were excluded from the study. The sample size was estimated as 30 patients per group by a power of 80%, effect size of 0.7 and confidence level of 95%. Treatment was started with tablet levothyroxine 0.1 mg (Iran Hormone Pharmaceutical Co, Tehran, Iran) with dose of 4  $\mu/kg/day$  before breakfast. Echocardiography was performed before and 6 months after starting levothyroxine treatment by the same cardiologist. The cardiologist was unaware of the intervention groups. Thyroid function tests (TSH and T4) were reassess 1.5 months after starting treatment and then every 2 months.

The obtained data were analyzed using SPSS statistical software. The descriptive analysis was used for the determination of the frequency, percentages, mean and standard deviation. Normality of the data was tested using the Kolmogorov-Smirnov test. Paired t-test and Wilcoxon signed rank were used to compare the parametric and non-parametric data, respectively, before and after the treatment. We utilized the exact method of bootstrap to calculate the p-value. A p-value of <0.05 was considered to be significant in all tests. The informed written consent was signed by all subjects' parents at the beginning of the study. The patients' information was kept confidential. This study was approved by the Ethics Research Committee of Babol University of Medical Sciences (code: IR.MUBABOL.HRI.REC.1397.153).

This trial was registered in the Iranian Registry of Clinical Trials with the number IRCT20180228038900N3.

## **Results**

Out of 30 subjects, 19 (63.3%) were boys and the rest were girls. The mean serum level of T4 was  $8.42\pm2.01~\mu\text{g/dl}$  before treatment and  $8.38\pm2.22~\mu\text{g/dl}$  at six months of treatment (p=0.86). The mean serum level of TSH was  $8.77\pm1.47~\text{mIu/l}$  before treatment and  $3.36\pm2.17~\text{mIu/l}$  at six months of treatment (<0.001). Figure 1 indicates the CONSORT flow diagram. Table 1 also shows the baseline information of the patients.




Figure 1. CONSORT flow diagram

Table 1. Baseline information of children with subclinical hypothyroidism

Variable	Results
Sex (boy/girl), number	19/11
Age, mean±SD, years old	6.60±2.13
Body Mass Index, mean±SD, kg/m <sup>2</sup>	16.55±4.00
T4, mean±SD, μg/dl	$8.42\pm2.01$
TSH, mean±SD, mIu/l	8.77±1.47

Table 2 shows comparison of cardiac functional parameters before and six months after treatment with levothyroxine in the patients. The mean EF index was 63.13±3.01 before treatment, which increased to 69.07±4.50 after treatment (p<0.001). Also, the mean FS was 31.83±1.62 before treatment, which improved to 35.10±1.13 after treatment (p<0.001). The mean MPI was 0.28±0.02 before treatment, which increased to 0.33±0.03 at six months of treatment (p<0.001). On the other hand, no significant

difference was found in the mean E/E' before and after treatment. The mean LVEDD decreased from  $3.47\pm0.46$  before treatment to  $3.05\pm0.40$  after treatment (p<0.001), whereas LVESD did not significantly decrease after treatment compared with before treatment. Table 3 indicates comparison of cardiac functional parameters before and six

months after treatment with levothyroxine by sex. In both boys and girls, EF, FS and MPI significantly increased after treatment compared with before treatment. LVEDD also significantly decreased after treatment compared with before treatment in both sexes. E/E' parameter increased in boys, but did not significantly increase in girls.

Table 2.Cardiac functional parameters before and six months after treatment with levothyroxine, in children with subclinical hypothyroidism

70113 1 0 10125 111			
Variables	Before treatment (mean±SD)	Six months of treatment (mean±SD)	P-value
Systolic parameters			
$\mathrm{EF}^{\mathrm{a}}\left(\% ight)$	63.13±3.01	69.07±4.50	< 0.001
FS <sup>b</sup> (%)	$31.83\pm1.62$	35.10±1.13	< 0.001
Diastolic parameter			
E/E <sup>rc</sup>	6.31±0.71	$6.49\pm0.74$	0.07
Global ventricular performance			
$\mathrm{MPI}^{\mathrm{d}}$	$0.28\pm0.02$	$0.33\pm0.03$	< 0.001
Cardiac morphological parameters			
LVEDD <sup>e</sup> (cm)	3.47±0.46	$3.05\pm0.40$	< 0.001
LVESD <sup>f</sup> (cm)	2.08±0.36	2.02±0.44	0.64

a Ejection fraction; b Fractional shortening; c Ratio of early filling velocity to early diastolic mitral annulus velocity; d myocardial performance index; e left ventricular end-diastolic diameter; f left ventricular end-systolic diameter

Table 3. Cardiac functional parameters before and six months after treatment with levothyroxine, in children with subclinical hypothyroidism by sex

Sex	Variables	Before treatment (mean±SD)	Six months of treatment (mean±SD)	P-value
Boys	Systolic parameters			
	EF <sup>a</sup> (%)	62.95±3.29	69.63±4.17	< 0.001
	FS <sup>b</sup> (%)	31.42±1.50	$35.05\pm2.19$	< 0.001
	Diastolic parameter			
	E/E <sup>c</sup>	$6.34\pm0.63$	$6.60\pm0.72$	0.010
	Global ventricular performance			
	$\mathbf{MPI}^{\mathrm{d}}$	$0.28\pm0.03$	$0.33\pm0.02$	< 0.001
	Cardiac morphological parameters			
	LVEDD <sup>e</sup> (cm)	$3.49\pm0.48$	$3.01\pm0.45$	< 0.001
	LVESD <sup>f</sup> (cm)	$2.09\pm0.30$	$2.00\pm0.48$	0.990
Girls	Systolic parameters			
	EF (%)	63.45±2.52	68.09±5.06	0.040
	FS (%)	32.55±1.63	35.18±2.13	0.010
	Diastolic parameter			
	E/E'	6.27±0.87	6.31±0.76	0.84
	Global ventricular performance			
	MPI	$0.28\pm0.01$	$0.33\pm0.04$	0.003
	Cardiac morphological parameters			
	LVEDD (cm)	$3.42\pm0.45$	3.12±0.30	0.006
	LVESD (cm)	2.06±0.17	2.05±0.39	0.380

a Ejection fraction; b Fractional shortening; c Ratio of early filling velocity to early diastolic mitral annulus velocity; d myocardial performance index; e left ventricular end-diastolic diameter; f left ventricular end-systolic diameter

## **Discussion**

To our knowledge, this is the first study in Iran that investigated the effects of levothyroxine on cardiac function in children with subclinical hypothyroidism using systolic, diastolic and morphologic parameters. As observed, systolic parameters improved after treatment. In addition, treatment with levothyroxine had positive effects on MPI and LVEDD.

So far, limited data has been published similar to our topic. In one study by Catlı et al. (27) in Turkey, which was performed on 31 children with subclinical hypothyroidism, it was seen that the morphological parameters of LVEDD and LVESD increased in 6 months of treatment that the first result was in agreement with our finding, but the latter was not compatible with our data. Çatlı et al. (27) also mentioned that other cardiac functional parameters, such as EF, FS and MPI did not significantly change after 6 months, which were contrary to our results. The authors stated that short duration of treatment might be a reason for lack of changes. The incompatibility between their results and ours might be due to differences in age, BMI, and/or serum levels of TSH and T4. Adult studies did not only showe that patients with subclinical hypothyroidism had impaired myocardial contractility and diastolic dysfunction, but also stated that these problems may be reversed with levothyroxine treatment (28-31). The study by Nakova et al. (10) on 54 patients showed that a 5-month levothyroxine treatment could contribute to higher EF and lower E/E'. Another research by Tadic et al. (32) on 54 Serbian women with subclinical hypothyroidism indicated that parameter of E/E' improved after 1-year treatment, but no changes were found in LVEDD and LVESD. The last results were compatible with the article by Ilic et al. (33), in which they reported that no significant changes were seen in LVEDD and LVESD after 1-year treatment with levothyroxine. A large-scale cohort study by Andersen et al. (34) revealed that levothyroxine treatment did not reduce risk of mortality in patients with subclinical hypothyroidism and heart disease. These data are helpful for clinicians to select a preventive and therapeutic approach against cardiac complications of subclinical hypothyroidism through levothyroxine administration, however, stronger evidence is needed in this regard. We also evaluated the effect of levothyroxine by sex. The notable point found from the results was improvement of E/E' parameter in boys but not in girls after treatment. This may be explained by differences in hormonal profiles between sexes. To the best of our knowledge, except Çatlı et al.' study, there are no other children studies to compare our findings with. Considering that the frequency of children with subclinical hypothyroidism is not low and they impose high economic costs on families and societies, addition of cardiac compactions following subclinical hypothyroidism increase pressure on children and their family. Therefore, more studies are suggested to be performed on children and adolescent patients with subclinical hypothyroidism. These studies can increase awareness and information about therapeutic drugs and help clinicians for better management of the patients.

In most of the papers, it has been exhibited that subclinical hypothyroidism is related to different cardiac functional parameters (35, 36). Studies on patients with subclinical hypothyroidism, who underwent levothyroxine treatment, showed evidence of improving left ventricular systolic function. Therefore, subclinical hypothyroidism can be considered as a mild hypothyroidism which is associated with primary cardiovascular symptoms. Thus, starting early and timely treatment in these patients seems necessary to prevent cardiac involvement.

One of the limitations of our study was the small sample size and relatively short follow-up period of study participants. Second, our study did not have an untreated control group (e.g., including patients who did not agree to undergo levothyroxine treatment) and we cannot certainly and strongly say that the observed improvements of cardiac function were a result of the levothyroxine treatment and not other factors, like physiological growth. Overall, it has been proposed to design new studies with bigger sample size, longer follow-up period and an untreated control group, plus assess other cardiac functional indices, such as left ventricle mass index, thickness of the left ventricle posterior wall, and mitral deceleration time.

In conclusion, according to the results, it can be concluded that treatment with levothyroxine may improve cardiac functional parameters in the children with subclinical hypothyroidism. However, it is necessary to perform new studies with an untreated control group to confirm the positive effect of levothyroxine on cardiac functions in childhood subclinical hypothyroidism.

## **Acknowledgments**

We are grateful to the Clinical Research Development Committee of Amirkola Children's Hospital, the Research Council of Non-Communicable Pediatric Diseases Research Center and Health Research Institute of Babol University of Medical Sciences for their support and cooperation in the study.

**Funding:** This study was supported by a research grant and residency thesis of Dr Nasrin Bano Rupani from the Non-Communicable Pediatric Diseases Research Center of Babol University of Medical Sciences (Grant Number: 9706119). **Conflict of interest:** None declared.

#### References

- Cojić M, Cvejanov-Kezunović L. Subclinical hypothyroidism - whether and when to start treatment? Open Access Maced J Med Sci 2017; 5: 1042-6.
- Such K, Gawlik A, Dejner A, et al. Evaluation of subclinical hypothyroidism in children and adolescents: a single-center study. Int J Endocrinol 2016; 2016: 1671820.
- 3. Tng EL. The debate on treating subclinical hypothyroidism. Singapore Med J 2016; 57: 539-45.
- Duntas LH, Chiovato L. Cardiovascular risk in patients with subclinical hypothyroidism. Eur Endocrinol 2014; 10: 157-60.
- Jabbar A, Ingoe L, Pearce S, Zaman A, Razvi S. Thyroxine in acute myocardial infarction (ThyrAMI)levothyroxine in subclinical hypothyroidism post-acute myocardial infarction: study protocol for a randomised controlled trial. Trials 2015; 16: 115.
- Fatourechi V. Subclinical hypothyroidism and subclinical hyperthyroidism. Expert Rev Endocrinol Metab 2010; 5: 359-73.
- 7. Floriani C, Gencer B, Collet TH, Rodondi N. Subclinical thyroid dysfunction and cardiovascular diseases: 2016 update. Eur Heart J 2017; 39: 503-7.
- 8. Suh S, Kim DK. Subclinical Hypothyroidism and Cardiovascular Disease. Endocrinol Metab (Seoul) 2015; 30: 246-51.
- 9. Biondi B, Klein I. Hypothyroidism as a risk factor for cardiovascular disease. Endocrine 2004; 24: 1-13.
- 10. Nakova VV, Krstevska B, Kostovska ES, Vaskova O, Ismail LG. The effect of levothyroxine treatment on left ventricular function in subclinical hypothyroidism. Arch Endocrinol Metab 2018; 62: 392-8.

- 11. Wiersinga WM. Guidance in subclinical hyperthyroidism and subclinical hypothyroidism: are we making progress? Eur Thyroid J 2015; 4: 143-8.
- 12. Kahaly GJ, Dillmann WH. Thyroid hormone action in the heart. Endocr Rev 2005; 26: 704-28.
- 13. Nicolini G, Pitto L, Kusmic C, Balzan S, et al. New insights into mechanisms of cardioprotection mediated by thyroid hormones. J Thyroid Res 2013; 2013: 264387.
- 14. Aksoy DY, Cinar N, Harmanci A, et al. Serum resistin and high sensitive CRP levels in patients with subclinical hypothyroidism before and after L-thyroxine therapy. Med Sci Monit 2013; 19: 210-5.
- 15. Mancini A, Di Segni C, Raimondo S, et al. Thyroid hormones, oxidative stress, and inflammation. Mediators Inflamm 2016; 2016: 6757154.
- 16. Hajje G, Saliba Y, Itani T, et al. Hypothyroidism and its rapid correction alter cardiac remodeling. PLoS One 2014; 9: e109753.
- 17. Shatynska-Mytsyk I, Rodrigo L, Cioccocioppo R, et al. The impact of thyroid hormone replacement therapy on left ventricular diastolic function in patients with subclinical hypothyroidism. J Endocrinol Invest 2016; 39: 709-13.
- 18. Martins RM, Fonseca RH, Duarte MM, et al. Impact of subclinical hypothyroidism treatment in systolic and diastolic cardiac function. Arq Bras Endocrinol Metabol 2011; 55: 460-7.
- 19. Razvi S, Weaver JU, Butler TJ, Pearce SH. Levothyroxine treatment of subclinical hypothyroidism, fatal and nonfatal cardiovascular events, and mortality. Arch Intern Med 2012; 172: 811-7.
- 20. Kalra P, Yeragani VK, Prasanna Kumar KM. Cardiac autonomic function and vascular profile in subclinical hypothyroidism: Increased beat-to-beat QT variability. Indian J Endocrinol Metab 2016; 20: 605-11.
- 21. Banker A, Bell C, Gupta-Malhotra M, Samuels J. Blood pressure percentile charts to identify high or low blood pressure in children. BMC Pediatr 2016; 16: 98.
- 22. Marshall WA, Tanner JM. Variations in the pattern of pubertal changes in boys. Arch Dis Child 1970; 45: 13-23.
- 23. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. Arch Dis Child 1969; 44: 291-303.
- 24. Bernstein D. History and physical exam. In: Kliegman RM, Stanton BF, St. Geme JW III, Schor NF, Behrman

- RE, eds. Nelson Textbook of Pediatrics. 20th ed. Philadelphia (PA), USA: Elsevier Saunders 2016; pp. 2163-2170.e1
- 25. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, et al. Recommendations for chamber quantification. Eur J Echocardiogr 2006;7: 79-108.
- 26. Bruch C, Klem I, Breithardt G, Wichter T, Gradaus R. Diagnostic usefulness and prognostic implications of the mitral E/E' ratio in patients with heart failure and severe secondary mitral regurgitation. Am J Cardiol 2007; 100: 860-5.
- 27. Çatlı G, Kır M, Anık A, et al. The effect of L-thyroxine treatment on left ventricular functions in children with subclinical hypothyroidism. Arch Dis Child 2015; 100: 130-7.
- 28. Turhan S, Tulunay C, Ozduman Cin M, et al. Effects of thyroxine therapy on right ventricular systolic and diastolic function in patients with subclinical hypothyroidism: a study by pulsed wave tissue Doppler imaging. J Clin Endocrinol Metab 2006; 91: 3490-3.
- 29. Erkan G, Erkan AF, Cemri M, et al. The evaluation of diastolic dysfunction with tissue Doppler echocardiography in women with subclinical hypothyroidism and the effect of L-thyroxine treatment on diastolic dysfunction: a pilot study. J Thyroid Res 2011; 2011: 654304.
- 30. Monzani F, Di Bello V, Caraccio N, et al. Effect of levothyroxine on cardiac function and structure in

- subclinical hypothyroidism: a double blind, placebocontrolled study. J Clin Endocrinol Metab 2001; 86: 1110-5.
- 31. Franzoni F, Galetta F, Fallahi P, et al. Effect of L-thyroxine treatment on left ventricular function in subclinical hypothyroidism. Biomed Pharmacother 2006; 60: 431-6.
- 32. Tadic M, Ilic S, Kostic N, Caparevic Z, Celic V. Subclinical hypothyroidism and left ventricular mechanics: a three-dimensional speckle tracking study. J Clin Endocrinol Metab 2014; 99: 307-14.
- 33. Ilic S, Tadic M, Ivanovic B, et al. Left and right ventricular structure and function in subclinical hypothyroidism: The effects of one-year levothyroxine treatment. Med Sci Monit 2013; 19: 960-8.
- 34. Andersen MN, Olsen AS, Madsen JC, et al. Long-term outcome in levothyroxine treated patients with subclinical hypothyroidism and concomitant heart disease. J Clin Endocrinol Metab 2016; 101: 4170-7.
- 35. Brienza C, Grandone A, Di Salvo G, et al. Subclinical hypothyroidism and myocardial function in obese children. Nutr Metab Cardiovasc Dis 2013; 23: 898-902.
- 36. Chen X, Zhang N, Cai Y, Shi J. Evaluation of left ventricular diastolic function using tissue Doppler echocardiography and conventional doppler echocardiography in patients with subclinical hypothyroidism aged< 60 years: a meta-analysis. J Cardiol 2013; 61: 8-15.